

CHEATHAM (W.)

Ophthalmia Neonatorum, by William Cheatham, M. D., Professor of Ophthalmology, Otology and Laryngology in the Louisville Medical College, Louisville, Ky.







OPHTHALMIA NEONATORUM.


By WILLIAM CHEATHAM, M.D.,

Professor of Ophthalmology, Otology and Laryngology in the Louisville Medical College, Louisville, Ky.

N this short paper on Ophthalmia Neonatorum I will give only the treatment that has given me the most satisfaction: On first seeing a case of ophthalmia neonatorum, I warn all the attendants of the danger to their own eves as well as to those of the patient. Some years ago I saw a grandmother lose both eyes as the result of inoculation from her grandchild, not having been warned of this danger, although the patient made a perfect recovery. On first seeing the case, [order (1) a saturated solution of boric acid, for frequent cleansing; (2) glyceride of tannin, 3 i, aqua dest. q. s., \(\frac{7}{3} \) i; (3) atropia sulph. gr. ss., saturated solution acid boric, \(\frac{7}{3}\) ss., and (4) argent, nitrate gr. ii, gr. iv, or gr. vi, to aqua dest., 3 ss. Four droppers are ordered, and the nurse is directed to put on one bottle a red or some colored string, with a corresponding color on the dropper used in that medicine, so the droppers and medicine cannot become mixed. Acid boric gr. xv, vaseline \(\frac{7}{2} \) i, is ordered to be used on the lids to prevent glueing and to protect the skin during the frequent bathing. The bottles are numbered 1, 2, 3, 4 and 5. No. 1 is the boric acid, No. 2 tannic acid, No. 3 atropia, No. 4 argent, nitrate, No. 5 salt water. The following directions are given, the numbers on the bottles only used: Bathe eyes well with No. 1 (boric acid solution) every hour; four times a day, after No. 1 has been used, drop into each eye three or four drops of No. 2 (glyceride of tannin); this cleanses the cul de sacs well; the lids should be manipulated thoroughly, and the secretions will come out in coagulated threads; the eyes are to be washed out with No. 1 (boric acid solution again); night and morning No. 3 (atropia solution) is to be dropped in: I do not wait for involvement of cornea and iris. Four times a day No. 4 (argent, nitrate solution) is used after No. 2 (glyceride of tannin) is used, and the lids manipulated well, so as to reach the cul de sacs; immediately afterwards, the eyes are flushed well with sodium chloride 3 i, aqua Oi. Then the lids are well covered with the ointment.

The glyceride of tannin serves a double purpose; by coagulating the secretion it acts as a cleanser, and then its action on the inflamed mucous membrane is favorable. Hydrogen dioxide is





sometimes used for the same purpose; I think its use dangerous, as it seems to have an unfavorable action upon the corneal epithelium, softening it and producing abrasions, which are followed by sloughs, loss of vision, etc.

I direct that nothing be used about the eye for cleansing, such as sponges, but instead old linen and clean surgical cotton, or such material as can be burned. When there is much edema of the lids, hot or cold applications are of much service. Recently I have had most excellent results from the use of Formol 1 to 4,000 or 1 to 5,000 sol. In ulceration of the cornea, Formol gives most wonderful results.

When these directions are followed from the first day of the disease, I believe no eyes will be lost.

Published on the 1st and 15th of the Month, \$2 or 8 Shillings a Year.

An Illustrated Semi-Monthly Devoted to the

DISEASES OF CHILDREN.

DILLON BROWN, M. D. NEW YCRK.

GEO. A. CARPENTER, M. D. LONDON.

st st st

EDITORIAL STAFF:

Medicine.

A. Jacobi, M. D., New York, Clinical Professor of Diseases of Children in Columbia College.

Frederick Forchheimer, M.D., Cincinnati, Ex-President of the American Pediatric Society, and Professor of Diseases of Children in the Medical College of Ohio.

Surgery.

Henry R. Wharton, M. D., Philadelphia, Surgeon to the Children's and the Presbyterian Hospitals. Frederic S. Eve, F. R. C. S., Eng., London, Surgeon to the London Hospital and to the Evelina Hospital

Lambert H. Ormsby, M. D., F. R. C. S. I., Dublin, Surgeon Meath Hospital and National Children's

Orthopedics.

Henry Ling Taylor, M. D., New York, Assistant to the Hospital for Ruptured and Crippled. Frederic R. Fisher, F. R. C. S., Eng., London, Senior Surgeon to the National Orthopedic Hospital, and late Surgeon to the Victoria Hospital for Children.

Therapeutics.

Reynold Webb Wilcox, M. D., New York, Professor of Therapeutics in the Post-Graduate Medical School, and Physician to St. Mark's Hospital.

John Thomson, M. D., Edinburgh, Extra Physician Royal Hospital for Sick Children, and Lecturer on Diseases of Children, Edinburgh School of Medicine.

Dawson Williams, M. D., London, Physician to the East London Hospital for Children.

Teratology.

Egbert H. Grandin, M. D., New York, Ex-President of the County Medical Society: Consulting Obstetric Surgeon to the Maternity Hospital, and Gynecologist to the French Hospital.

J. W. Ballantyne, M. D., Edinburgh, Lecturer on the Diseases of Infancy and Childhood, Minto House; and Physician for Diseases of Children, Convente Disbersary.

Cowgate Dispensary.

Gastro-Enteric Diseases.

Morris Manges, M. D. New York, Physician to Mount Sinai Hospital, and Editor of "Ewald's Diseases of the Stomach." J. Boas, M. D. Berlin, Editor of the "Archiv fur Verdanungs-krankheiten."

Genito-Urinary Diseases.

Frederic Russell Sturgis, M. D., New York, Visit-

ing Surgeon to the City Hospital.

Fancourt Barnes, M. D., London, Consulting Physician to the British Lying-In Hospital, S. George's Hospital, Seamen's Hospital and East London Hospital for Children.

Neurology.

Joseph Collins, M. D., New York, Physician to the Hospital for Nervous Diseases.

James Taylor, M. D., London, Senior Assistant Physician to the National Hospital for the Paralysed.

Materia Medica.

Henry H. Rusby, M. D., New York, Professor of Materia Medica and Botany in the College of Pharmacy.

Laryngology and Otology.

Wm. C. Glasgow, M. D., St. Louis, Professor of Laryngology in the Missouri Medical Coalege.

Laryngology in the Missouri Medical Coolege.

Arthur H. Cheatle, F. R. C. S., London, Assistant Surgeon to the Royal Ear Hospital and Lural Department of King's College Hospital.

Max Thorner, M. D., Cincinnati College of Medicine and Surgery; and Aurist and Laryngologist to the Cincinnati Hospital.

Peter McBride, M. D., Edinburgh, Aural Surgeon and Laryngologist, Royal Infirmary; and Lecturer on the Diseases of the Ear and Throat, Edinburgh School of Medicine.

Dermatology.

James Nevins Hyde, M.D., Chicago, Professor of Skin and Venereal Diseases in Rush Medical College. Leslie Phillips, M.D., Birmingham, England, Sur-geon Birmingham and Midland Skin and Lock Hospital.

Ophthalmology.

Myles Standish, M. D., Boston, Ophthalmic Surgeon to Carney Hospital and to Massachusetts Charity Eye and Ear Infirmary.

Eye and Ear Infirmary.
William Arthur Brailey, M. D., London, Ophthalmic Surgeon to Evelina Hospital for Children, and
Guy's Hospital.

Pathology and Bacteriology.

Henry Ashby, M. D., Manchester, England. Physician Manchester General Hospital for Children; and Lecturer on Diseases of Children, Owen's

Alfred Lingard, M. D., Poona, India, Professor of Bacteriology, Science College.

Physiology.

C. S. Sherrington, M. D., F. R. S., Liverpool, England, Holi Professor of Physiology, University College.

Psychology.

T. Telford-Smith, M. D., Lancaster, England, Resident Medical Superintendent, Royal Albert Asylum.

Tropical Diseases.

Surgeon-Major H. P. Birch, Quetta, India.

SEND FOR A SAMPLE COPY.

VOL.1 NO1

JANUARY 151 1896

\$2 OR 8s A YEAR

PEDIATRICS



VAN PUBLISHING CQ 1432 BROADWAY CQR 40TH ST, NEW YORK JOHN BALE & SONS, 85-89 GREAT TITCHFIELD ST,W, LONDON